L# ANSWER 75 OF 419 CAPLUS COPYRIGHT 2002 ACS

AN 1998:389150 CAPLUS

DN 129:67694

TI Preparation of N-methyl-2-pyrrolidone from .gamma.-butyrolactone and monomethylamine

IN Nishida, Mitsuhiro; Komatsu, Yshinori; Mukugi, Tomohiro; Nakamura, Teiji

PA Tonen Chemical Corp., Japan

SO Jpn. Kokai Tokkyo Koho, 4 pp.

CODEN: JKXXAF

PATENT NO. KIND DATE APPLICATION NO. DATE

------

PI JP 10158238 A2 19980616 JP 1996-334446 19961202

AB The title compd. is prepd. by reaction of .gamma.-butyrolactone with 1.03-1.50 mol equiv. MeNH2 in the presence of 1.0-2.9 mol equiv. H2O at 250-300.degree. .gamma.-Butyrolactone (3 mol) was treated with 3 mol H2O and 3.3 mol MeNH2 at 280.degree. for 1 h to give 99.9% N-methyl-2-pyrrolidone. An amine mixt. contg. MeNH2 39.2, Me2NH 0.67, and Me3N 0.35 wt.% was recovered from the reaction mixt. and recycled to the reaction.

IC ICM C07D207-267

DT Patent

LA Japanese